

RICEBEAUX® HERBICIDE

FOR DISTRIBUTION AND USE ONLY IN ARKANSAS
SUPPLEMENTAL LABEL

TO BE USED IN CONJUNCTION WITH COMMAND 3ME SLN LABEL

EPA REG. NO.: 71085-30

FOR AERIAL APPLICATION ONLY OF RICEBEAUX® HERBICIDE IN A THREE WAY TANK-MIX WITH
COMMAND® 3ME AND ONE OF THE FOLLOWING PRODUCTS ON RICE IN THE STATE OF ARKANSAS:

PERMIT PLUS, FACET 75 DF, NEWPATH, PROWL H20 AND FACET L

OR A TWO WAY TANK-MIX WITH RICEBEAUX AND COMMAND.

FOR AGRICULTURAL OR COMMERCIAL USE ONLY

THIS LABEL FOR RICEBEAUX HERBICIDE IS VALID UNTIL WITHDRAWN, CANCELLED OR SUSPENDED.

ALL APPLICABLE DIRECTIONS, RESTRICTIONS AND PRECAUTIONS ON ALL EPA REGISTERED LABELS
AND COMMAND 3ME SLN No. AR08-0001 LABELS ARE TO BE FOLLOWED.

THIS LABEL MUST BE IN THE POSSESSION OF THE USER AT THE TIME OF PESTICIDE APPLICATION.

IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH ITS
LABELING.

DIRECTIONS FOR USE:

For aerial application only:

WEEDS CONTROLLED

Barnyardgrass (watergrass)
Broadleaf signalgrass
Coffeebean
Coffeeweed
Crabgrass
Croton
Dayflower
Eclipta
False Pimpernel
Flatsedge
Foxtail
Goosegrass
Gulf cockspur
Hemp Sesbania
Hoorahgrass
Indigo
Jointvetch, Northern and Indian
Junglerice
Mexicanweed
Millet (Texas)
Morningglory, Pitted

Echinochloa crus-galli
Brachiaria platyphylla.
Hemp Sesbania
Sesbania herbacea
Digitaria spp.
Croton spp.
Commelina communis
Eclipta prostrata
Lindernia spp.
Cyperus erythrorhizos, C. iria
Setaria spp.
Eleusine indica
Echinochloa crus-pavonis
Sesbania exaltata
Fimbristylis spp.
Aeschynomene virginica
Aeschynomene spp.
Echinochloa colonum
Caperonia castanifolia
Urochloa texana
Ipomoea lacunosa

Panicum, Fall
Paragrass
Pigweed
Redstem
Sicklepod
Sourdock, Curly Dock
Spearhead
Spikerush
Waterhyssop
Wiregrass
Sprangletop
Small ducksalad*
Smartweed
Yellow nutsedge

Panicum dichotomiflorum
Urochloa mutica
Amaranthus spp.
Ammannia spp.
Cassia obtusifolius
Rumex crispus
Phacelia hastata
Eleocharis obtuse, E. parvula
Bacopa rotundifolia
Eleusine indica
Leptochloa spp.
Heteranthera spp.
Polygonum spp.
Cyperus esculentus

*prior to the spoon leaf stage

(RiceBeaux will not control arrowhead, Bermudagrass, cattail, ducksalad, Johnsongrass, nutgrass and red rice,). REFER TO THE COMMAND 3ME AND OTHER PRODUCT LABELS FOR ADDITIONAL WEEDS CONTROLLED/SUPPRESSED.

Apply 3 quarts RiceBeaux with Command® 3 ME and selected tank mix product as a three way tank-mix or RiceBeaux and Command 3ME as a 2 way tank mix. This tank-mix should be applied after rice emerges (1-2 leaf stage) for control of existing weeds present at the time of aerial application.

Observe all buffer restrictions noted under Product Restrictions and Application Restrictions/ Precautions sections.

CROP SAFETY PRECAUTIONS:

Application to fields which have been precision leveled with deep cuts may result in rice crop injury including stand loss. Consult with rice specialists for soil amending practices which can reduce potential for herbicide injury in precision leveled fields.

AERIAL APPLICATION REQUIREMENTS:

Refer to Arkansas Pesticide Use and Application Act and Regulations. Act 389 or 1975 as amended (pages 15-17).

APPLICATION PRECAUTIONS:

Caution must be taken to minimize off target drift of this tank-mix as off-site movement can cause foliar whitening or yellowing of some plants.

- Do not apply with 1200 feet of the following areas:
- Towns and Housing Developments
- Commercial Vegetable Production (except peppers, pumpkins, succulent peas, sweet corn, sweet potato and winter squash)
- Commercial Fruit/Nut Production
- Commercial Nurseries
- Commercial Greenhouses
- Avoid making applications of the tank-mix when spray particles may be carried by air currents to areas where sensitive crops and plants are growing, or when temperature inversions exist. Prior to application, adjacent properties must be checked, and applications within 300 feet of desirable plants must be avoided.

RICEBEAUX PRODUCT RESTRICTIONS:

IN ARKANSAS: The following use prohibitions apply in Cross, Lee, Mississippi, Poinsett and St Frances Counties:

1. **RiceBeaux®** is not to be applied aurally within one mile of the St. Francis Floodway (west branch of St. Francis River) where the fat pocket book pearly mussel is known to occur;
2. **RiceBeaux** is not to be ground applied within 1,000 feet of the St. Francis Floodway where the fat pocket book pearly mussel is known to occur.
3. Rice fields are not to be flooded for at least 3 days after application, and water application on the fields is not to be drained for at least 7 days after flooding a treated field in areas where waters drain into the St. Francis Floodway where the fat pocket book pearly mussel if known to occur; and
4. Should on-going distributional surveys of the fat pocketbook pearly mussel find additional populations in the St. Francis Floodway, or other waters, the same restrictions would apply to these waters.

DO NOT apply this product south of the Intracoastal Waterway in Louisiana

DO NOT apply this product within two (2) miles from the shorelines of Matagorda Bay in Texas.

DO NOT apply this product within two (2) miles from the shorelines of Galveston Bay in Texas.

DO NOT plant or transplant crops in the treated area for 60 days following an application of this product.

DO NOT apply more than 5.3 qts. **RiceBeaux** per acre per application.

DO NOT apply more than 5.3 qts. **RiceBeaux** (4 lbs active ingredient propanil/4 lbs active ingredient thiobencarb) per season.

Applications to fields where catfish/crayfish farming is practiced and draining water from treated fields into areas where catfish farming is practiced is prohibited during 12 months following treatment. Do not use adjacent to catfish/crayfish ponds.

In Texas and other areas where double cropping is the agricultural practice, do not apply to a second stubble rice crop.

DO NOT apply this product in a way that will contact workers or other persons either directly or through drift. Only protected handlers may be in the area during application.

DO NOT apply this product (directly or indirectly) to any crop except rice.

DO NOT release permanent flood water within 14 days of application of this product (where weather permits).

Avoid application of this product within 24 hours of rainfall, or when heavy rain is expected to occur within 24 hours.

DO NOT apply when wind conditions will allow drift to adjacent, susceptible crops such as beans, soybeans, cotton, safflower, cucurbits, vegetables, orchards and other sensitive crops.

DO NOT harvest within 60 days of application.

Water drained from treated rice fields must not be used to irrigate other crops or be released within ½ miles upstream of a potable water intake in flowing water (i.e., river, stream, etc.) or within ½ miles of a potable water intake in a standing body of water such as a lake, pond, or reservoir.

DO NOT apply with 14 days before or after organophosphate or carbamate insecticide application.

DO NOT apply when rain is expected within 6 hours.

Rice seedlings with succulent growth may exhibit temporary foliar burn which may be greater than conventional Propanil application but usually recover after 10 to 14 days.

DO NOT mix liquid nitrogen, zinc or surfactants.

DO NOT exceed any label dosage rates.

DO NOT apply to second crop (stubble crop) rice.

DO NOT apply to fields with exposed seed as exposed seed will be killed.

DO NOT overlap or double spray ends of field.

DO NOT apply when temperature exceeds 90°F

DO NOT mix/load or otherwise handle this product within 100 feet of aquatic habitat.

This product cannot be mixed with any product containing a label prohibiting against such mixing.

Application to stressed rice can result in stand reduction, chlorosis, growth inhibition, delayed maturity and/or leaf desiccation. Stress factors include but are not limited to the following: Daily temperature below 65°F or above 95°F, problem soils, (i.e., Zn deficiency, high salt content, high pH), excessive moisture, (i.e., above field capacity while rice seed is germinating), drought conditions, poor field drainage or deep water after application.

Refer to product labels for Command 3ME and other tank-mix partners for additional restrictions.

When using Facet, or Facet L, refer to product label for additional Arkansas restrictions.

ROTATIONAL CROP GUIDELINES*:

Refer to Command 3ME and other tank mix partner federal labels for all rotational crop guidelines and restrictions.

*Refer to "REQUIREMENTS FOR PLANTING TIME APPLICATIONS" and "REPLANTING INSTRUCTIONS" in the Command 3ME and other tank mix partner federal labels.

Manufactured for:

RICECO LLC

5100 Poplar Avenue, Suite 2428

Memphis, TN 38137

1/7/13

RiceBeaux is a registered trademark of RiceCo LLC

Command is a registered trademark of FMC

Facet, Facet L, Prowl H20 and Newpath are registered trademarks of BASF

Permit Plus is a registered trademark of Gowan