SPECIAL LOCAL NEED 24(C)LABELING

DuPont Crop Protection

DUPONT™ LANNATE® SP INSECTICIDE SOYBEANS STATE OF ARKANSAS

RESTRICTED USE PESTICIDE

DUE TO HIGH ACUTE TOXICITY TO HUMANS.

For retail sale and use only by Certified Applicators or persons under their direct supervision and only for those uses covered by the Certified Applicator's certification. Direct supervision for this product requires the certified applicator to review federal and supplemental label instructions with all personnel prior to application, mixing, loading, or repair or cleaning of application equipment.

FOR DISTRIBUTION AND USE ONLY WITHIN THE STATE OF ARKANSAS SUPPLEMENTAL LABELING

DUPONTTM LANNATE® SP INSECTICIDE

EPA Reg. No. 352-342

FOR CONTROL OF SOYBEAN LOOPER ON SOYBEANS IN THE STATE OF ARKANSAS

KEEP OUT OF REACH OF CHILDREN

DANGER PELIGRO



POISON CONTAINS METHOMYL

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand this label, find someone to explain it to you in detail.)

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling. DuPont™ LANNATE® SP Insecticide is recommended for use on soybeans to control soybean looper in Arkansas.

IMPORTANT

BEFORE USING LANNATE® SP, READ AND FOLLOW ALL APPLICABLE DIRECTIONS, RESTRICTIONS AND PRECAUTIONS ON THE EPA-REGISTERED LABEL.

This bulletin contains new or supplemental instructions for use of this product which do not appear on the EPA-registered package label. Follow the instructions carefully.

This labeling must be in the possession of the user at the time of pesticide application.

HOW TO USE

Apply 1/2 to 1 pound LANNATE® SP per acre in sufficient water to give thorough coverage. Use the lower rates on small larvae and light infestations and higher rates on large larvae and severe infestations.

Reentry into treated areas is prohibited for 48 hours unless personal protective clothing and equipment specified on the LANNATE® SP product label for early reentry are worn.

Do not apply more than 1.5 lbs of LANNATE® SP per acre per crop.

Do not make more than 3 applications a season per crop.

Do not apply within 14 days of harvest.

Do not allow livestock grazing/feeding on forage within 10 days and hay within 12 days of an application.

© 2005 E. I. du Pont de Nemours and Company, Crop Protection, Wilmington, Delaware 19898



© 2005 E. I. du Pont de Nemours and Company, Crop Protection, Wilmington, Delaware 19898